

## MA-231 Topology

### 3. Ordered Sets

August 20, 2004 ; Submit solutions **before 10:00 AM ; August 30, 2004.**

**3.1.** Let  $(X, \leq)$  be an ordered set in which every subset has a least upper bound and has a greatest lower bound. Further, let  $f : X \rightarrow X$  be an increasing map and let  $F$  be the fixed points of  $f$ . Show that:

- a). If  $\{x \in X \mid f(x) < x\} \neq \emptyset$  and if  $a$  is its greatest lower bound, then either  $a \in F$  or  $f(a) \in F$ .
- b). If  $\{x \in X \mid x < f(x)\} \neq \emptyset$  and if  $z$  is its least upper bound, then either  $z \in F$  or  $f(z) \in F$ .
- c).  $F$  is non-empty. Further, the least upper bound and the greatest lower bound of  $F$  belong to  $F$ .

**3.2.** Let  $(X, \leq)$  be a lattice (see T3.2) with a largest element and a smallest element.

a). Suppose that  $X$  has the following property : if  $x, y \in X$  and if  $\text{Sup}\{x, y\}$  is a *direct successor*<sup>1)</sup> of  $x$ , then  $\text{Inf}\{x, y\}$  is a *direct predecessor* of  $y$ . Prove the following Chain theorem: *If  $X$  has a finite maximal chain, then every chain in  $X$  is finite and the lengths<sup>2)</sup> of all maximal chains in  $X$  are equal.* (Hint: It is enough to prove that: if  $X$  has a finite maximal chain  $x_0 < \dots < x_n$  of length  $n$ , then every finite chain  $y_0 < \dots < y_m$  in  $X$  has length  $m \leq n$ . Induction on  $n$ : if  $n \geq 1, m \geq 1$ , then apply induction hypothesis on the lattice  $\{x \in M : x \leq x_{n-1}\}$ . In the case  $y_{m-1} \not\leq x_{n-1}$  consider the element  $\text{Inf}\{y_{m-1}, x_{n-1}\}$ . This element is  $\leq x_{n-1}$  and is a direct predecessor of  $y_{m-1}$ . If  $y_{m-2} \not\leq \text{Inf}\{y_{m-1}, x_{n-1}\}$ , then consider  $\text{Inf}\{y_{m-2}, \text{Inf}\{y_{m-1}, x_{n-1}\}\}$  and so on... (Induction on  $m$ ). Note that  $x_0$  resp.  $x_n$  is the smallest resp. greatest element in  $X$ .) — Give an example of a (with 5 elements) in which there are maximal chains of different lengths.

b). (Dedekind’s Chain Theorem) Suppose that  $X$  is artinian and noetherian and that  $X$  has the following property: if  $x, y \in X$  and if  $\text{Sup}\{x, y\}$  is a direct successor of  $x$  and  $y$ , then  $\text{Inf}\{x, y\}$  is a direct predecessor of  $x$  and  $y$ . Show that: All maximal chains in  $X$  have the same (finite) lengths. (Hint: Similar to that of part a).)

**3.3. a).** (Dilworth’s Theorem) Let  $(X, \leq)$  be a finite ordered set and let  $m$  be the cardinality of a largest possible *anti-chain*<sup>3)</sup> in  $X$ . Show that  $X$  can be partitioned into  $m$  chains and  $X$  cannot be partitioned into  $r$  chains with  $r < m$ . — This natural number  $m$  is called the Dilworth’s number of  $X$ . (Proof. By induction on the cardinality of  $X$ . Let  $Y \subseteq X$  be an anti-chain in  $X$  of cardinality  $m$ . Let  $A := \{a \in X \mid a < y \text{ for some } y \in Y\}$  and let  $B := \{b \in X \mid y < b \text{ for some } y \in Y\}$ . Then  $X = A \uplus B \uplus Y$ . We divide the proof in the following four cases. i).  $A = \emptyset = B$ . ii).  $A \neq \emptyset, B \neq \emptyset$ . iii).  $A \neq \emptyset, B = \emptyset$ . iv).  $A = \emptyset, B \neq \emptyset$ . The proof in the case i) is trivial. For case ii): Put  $X_A := Y \cup A$  and  $X_B := Y \cup B$ . Then by induction both  $X_A$  and  $X_B$  can be partitioned into  $m$  chains. Then  $X$  can also be partitioned into  $m$  chains. For the case iii): Let  $y \in Y$  and let  $C$  be a chain of maximal length with  $y \in C$ . Let  $X' := X \setminus C$ . Note that the extremal elements of  $C$  are also extremal elements of  $X$ . Let  $m'$  denote the the maximal number of pairwise incomparable elements in  $X'$ . Then  $m - 1 \leq m' \leq m$ , since  $Y \setminus \{y\} \subseteq X'$ . Choose an anti-chain  $Y'$  in  $X'$  with  $\text{card}(Y') = m'$ . We now further consider the following two cases: iii.a):  $m' = m$ . In this case replace  $Y$  by  $Y'$  and then apply the case ii) to complete the proof. iii.b):  $m' = m - 1$ . In this case apply induction hypothesis to  $X'$  to complete the proof. Proof in the case iv) is similar to that of case iii).)

b). (E. Sperner) Let  $X$  be a finite set with  $n$  elements. Show that the Dilworth’s number of the ordered set  $(\mathfrak{P}(X), \subseteq)$  is  $\binom{n}{\lfloor n/2 \rfloor}$ , where  $\lfloor n/2 \rfloor$  is the integral part of  $n/2$ . (Hint: Use the maps

$f_i, 0 \leq i < n/2$  and  $g_i, n/2 < i \leq n$  of Exercise 2.1 to give an explicit partition of  $\mathfrak{P}(X)$  into  $\binom{n}{\lfloor n/2 \rfloor}$  chains.

**Variant:** if  $\mathfrak{S} \subseteq \mathfrak{P}(X)$  be an anti-chain in  $\mathfrak{P}(X)$  then  $|\mathfrak{S}| \leq \binom{n}{\lfloor n/2 \rfloor}$  as follows: For  $Y \in \mathfrak{S}$ , let  $\mathfrak{C}_Y$  be the set of all maximal chains in  $\mathfrak{P}(X)$  in which  $Y$  appears as an element. Then  $|\mathfrak{C}_Y| = (n - |Y|)! \cdot |Y|!$  and  $\mathfrak{C}_Y \cap \mathfrak{C}_Z = \emptyset$  if  $Y, Z \in \mathfrak{S}, Y \neq Z$ . Since there are  $n!$  maximal chains, it follows that  $\sum_{Y \in \mathfrak{S}} (n - |Y|)! \cdot |Y|! \leq n!$  and so  $|\mathfrak{S}| \leq \binom{n}{\lfloor n/2 \rfloor}$ .) For  $1 \leq i \leq n$ , what is the Dilworth’s number of  $\mathfrak{P}_{\leq i}(X) := \{Y \in \mathfrak{P}(X) \mid |Y| \leq i\}$ ?

On the other side one can see (simple) test-exercises ; their solutions need not be submitted.

<sup>1)</sup> Let  $a, b$  be two elements in an ordered set  $(X, \leq)$ . If  $a < b$  and  $(a, b) := \{x \in X \mid a < x < b\} = \emptyset$ , then we say that  $b$  is a *direct successor* of  $a$  and we say that  $a$  is a *direct predecessor* of  $b$ .

<sup>2)</sup> By the length of a finite chain  $C$  in an ordered set  $(X, \leq)$ , we mean  $|C| - 1$ .

<sup>3)</sup> An anti-chain in an ordered set  $(X, \leq)$  is a subset of  $X$  consisting of pairwise incomparable elements.

### Test-Exercises

**T3.1. a).** Let  $(X, \leq)$  be a finite non-empty ordered set. Show that  $X$  has (at least one) a minimal and (at least one) a maximal element.

**b).** Let  $(X, \leq)$  be a finite totally ordered set with  $n$  elements. Show that there exists exactly one isomorphism of the interval  $[1, n] = \{1, \dots, n\} \subseteq \mathbb{N}$  (with the natural order) onto  $X$ .

**c).** Let  $(X, \leq)$  be an inductively ordered set and let  $x \in X$ . Show that there is a maximal element  $z \in X$  such that  $x \leq z$ . (Apply Zorn's <sup>†</sup> lemma <sup>4</sup>) to the subset  $\{y \in X \mid x \leq y\}$ .)

**T3.2. (Lattice)** An ordered set  $(X, \leq)$  is called a lattice if for every two elements  $x, y$  in  $X$ ,  $\text{Sup}\{x, y\}$  and  $\text{Inf}\{x, y\}$  exist. The power set of a set with respect to the natural inclusion is a lattice. Every totally ordered set is a lattice. If  $X_i, i \in I$ , is a family of lattices, then so is the product set  $\prod_{i \in I} X_i$  with respect to the product order. If  $X$  is a lattice, then  $X$  is also lattice with respect to the inverse order; this lattice is called the (dual lattice).

**T3.3.** Let  $X$  be a set. Then the power set  $\mathfrak{P}(X)$  of  $X$  is (with respect to the natural inclusion) noetherian resp. artinian if and only if  $X$  is finite.

**T3.4.** Let  $(X, \leq)$  be a well-ordered set.

**a).** Every element in  $X$  which is not the largest element in  $X$  has a direct successor in  $M$ . Is it true that every element in  $X$  which is not the smallest element in  $X$  necessarily have a direct predecessor in  $X$ ?

**b).** Let  $g$  be strictly increasing map from  $X$  into itself. Show that:  $x \leq g(x)$  for all  $x \in X$ . If  $\text{im } g$  is a section of  $X$ , then  $g = \text{id}_X$ . In particular,  $\text{id}_X$  is the only isomorphism of  $X$  onto itself. Further, if  $X, Y$  are well-ordered subsets, then there is at most one isomorphism of  $X$  onto a section of  $Y$ .

**T3.5.** Let  $(X, \leq)$  be an ordered set. A section of  $X$  is a subset  $A$  of  $X$  with the following property: if  $x \in A, y \in X$  with  $y \leq x$ , then  $y \in A$ . — The subsets  $\emptyset$  and  $X$  are sections of  $X$ . Arbitrary intersections and arbitrary unions of sections of  $X$  are again sections of  $X$ . For  $a \in X$ , the subsets  $A_a := \{x \in X \mid x < a\}$  and  $\bar{A}_a := \{x \in X \mid x \leq a\}$  are sections of  $X$ .

**a).** The map  $a \mapsto \bar{A}_a$  from  $X$  into  $\mathfrak{P}(X)$  is a strictly monotone increasing (where  $\mathfrak{P}(X)$  is ordered by the natural inclusion) and induces an isomorphism of  $X$  onto a subset of  $\mathfrak{P}(X)$ .

**b).** Suppose further that  $(X, \leq)$  is well-ordered. Then show that: If  $A$  is a section of  $X, A \neq X$ , then there exists exactly one  $a \in X$  such that  $A = A_a$ . The map  $a \mapsto A_a$  is an isomorphism of  $X$  onto the set of sections different from  $X$  which is ordered by the natural inclusion. The set of sections of  $X$  is well-ordered and has a greatest element.

<sup>†</sup> **Max August Zorn (1906-1993)** Max Zorn was born on 6 June 1906 in Krefeld, Germany and died on 9 March 1993 in Bloomington, Indiana, USA. Max Zorn was born in Krefeld in western Germany, about 20 km northwest of Dusseldorf. He attended Hamburg University where he studied under Artin. Hamburg was Artin's first academic appointment and Zorn became his second doctoral student. He received his Ph.D. from Hamburg in April 1930 for a thesis on alternative algebras. His achievements were considered outstanding by the University of Hamburg and he was awarded a university prize. He was appointed as an assistant at Halle but he did not have the opportunity to work there for long since, in 1933, he was forced to leave Germany because of the Nazi policies. He was not, however, Jewish. Zorn emigrated to the United States and was appointed a Sterling Fellow at Yale University. He worked there from 1934 to 1936 and it was during this period that he proposed "Zorn's Lemma" for which he is best known. Since Zorn is best known for "Zorn's Lemma" it is perhaps appropriate that we should begin a discussion of his mathematical achievements by considering this contribution. Of course Zorn did not call his result "Zorn's Lemma", rather it was given by him as a "maximum principle" in a short paper entitled A remark on method in transfinite algebra which he published in the Bulletin of the American Mathematical Society in 1935. Perhaps in passing we should note that the name "Zorn's Lemma" was due to John Tukey. Zorn's aim in this paper was to study field theory and in particular to improve on the method used for obtaining results in the subject. Methods used up to that time had depended heavily on the well-ordering principle which Zermelo had proposed in 1904, namely that every set can be well-ordered. What Zorn proposed in the 1935 paper was to develop field theory from the standard axioms of set theory, together with his maximum principle rather than Zermelo's well-ordering principle.

The form in which Zorn stated his maximum principle was as follows. The principle involved chains of sets. A chain is a collection of sets with the property that for any two sets in the chain, one of the two sets is a subset of the other. Zorn defined a collection of sets to be closed if the union of every chain is in the collection. His maximum principle asserted that if a collection of sets is closed, then it must contain a maximal member, that is a set which is not a proper subset of some other in the collection. The paper then indicated how the maximum principle could be used to prove the standard field theory results.

Today we know that the Axiom of Choice, the well-ordering principle, and Zorn's Lemma (the name now given to Zorn's maximum principle by Tukey and now the standard name) are equivalent. Did Zorn know this when he wrote his 1935 paper? Well at the end of the 1935 paper he did say that these three are all equivalent and promised a proof in a future paper. Was Zorn's idea entirely new? Well similar maximum principles had been proposed earlier in different contexts by several mathematicians, for example Hausdorff, Kuratowski and Brouwer. Paul Campbell. Following his years at Yale, he moved to the University of California at Los Angeles where he remained until 1946. During this time Herstein was one of his doctoral students. He left the University of California to become professor at Indiana University, holding this position from 1946 until he retired in 1971. After 1947 Zorn stopped publishing mathematical papers. This does not mean that he gave up mathematics. In recent years Max became fascinated by the Riemann Hypothesis and possible proofs using techniques from functional analysis. He read and studied and talked about mathematics nearly every day of his life. From time to time he published a slim newsletter. He was a gentle man with a sharp wit who, during nearly half a century, inspired and charmed his colleagues at Indiana University.

Max Zorn married Alice Schlottau and they had one son Jens and one daughter Liz.

<sup>4</sup> **Zorn's Lemma** Let  $(X, \leq)$  be an inductively ordered set, i.e. every chain in  $X$  has an upper bound in  $X$ . Then  $X$  has (at least one) a maximal element.