

HW 3 (to be submitted by 7 Feb)

1. (5 marks) Let $U \subset \mathbb{R}^n$ be an open set and $f : U \rightarrow \mathbb{R}^k$ (where $k \leq n$) be a smooth function. Suppose Df_a has rank k whenever $f(a) = 0$. Prove that $f^{-1}(0)$ can be made into a smooth manifold of dimension $n - k$.
2. (15 marks)
 - (a) (5 marks) Consider the complex projective space $\mathbb{C}\mathbb{P}^n = (\mathbb{C}^{n+1} - \{0\}) / (X \sim \lambda X \mid \lambda \in \mathbb{C} - \{0\})$. Give it a $2n$ -dimensional smooth manifold structure akin to that of $\mathbb{R}\mathbb{P}^n$.
 - (b) (5 marks) Prove that $(S^{2n+1} \subset \mathbb{C}^{n+1}) / (p \sim e^{i\theta} p)$ can be made into a $2n$ -dimensional smooth manifold such that the projection map from S^{2n+1} is smooth.
 - (c) (5 marks) Prove that $\mathbb{C}\mathbb{P}^n$ is diffeomorphic to $S^{2n+1} / (p \sim e^{i\theta} p)$. (Note that technically this problem is not well-defined because if you come up with crazy smooth structures on both spaces, then they are not diffeomorphic, but I highly doubt you can actually come up with non-diffeomorphic smooth structures!)
3. (5 marks) Let M be a smooth non-empty n -manifold ($n \geq 1$) with or without boundary. Prove that the space of smooth functions $f : M \rightarrow \mathbb{R}$ is infinite-dimensional.