

Finite-dimensional generalized nil-Coxeter and nil-Temperley–Lieb algebras

Apoorva Khare
Indian Institute of Science

December 13, 2024 @ ICTS Bangalore

Coxeter groups

The group S_{n+1} of permutations, and $(\mathbb{Z}/2\mathbb{Z}) \wr S_n$ of signed permutations, are examples of *finite Coxeter groups* – finite groups of orthogonal transformations generated by reflections.

- For S_{n+1} , let $s_1 = (1\ 2)$, $s_2 = (2\ 3)$, \dots , $s_n = (n\ n+1)$.

The **permutation group** S_{n+1} is generated by s_1, \dots, s_n with the **braid relations**

$$s_i s_j = s_j s_i \text{ (i.e. } (s_i s_j)^2 = 1), \quad |i - j| > 1, \quad s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1},$$

and the **Coxeter relations** $s_i^2 = 1$.

Coxeter groups

The group S_{n+1} of permutations, and $(\mathbb{Z}/2\mathbb{Z}) \wr S_n$ of signed permutations, are examples of *finite Coxeter groups* – finite groups of orthogonal transformations generated by reflections.

- For S_{n+1} , let $s_1 = (1\ 2)$, $s_2 = (2\ 3)$, \dots , $s_n = (n\ n+1)$.

The **permutation group** S_{n+1} is generated by s_1, \dots, s_n with the **braid relations**

$$s_i s_j = s_j s_i \text{ (i.e. } (s_i s_j)^2 = 1), \quad |i - j| > 1, \quad s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1},$$

and the **Coxeter relations** $s_i^2 = 1$.

- The **dihedral group** $I_2(m)$ (rotation and reflection symmetries of the regular m -gon) is generated by s_1, s_2 with the braid and Coxeter relations

$$(s_1 s_2)^m = 1 \text{ (i.e. } s_1 s_2 s_1 \cdots = s_2 s_1 s_2 \cdots), \quad s_1^2 = s_2^2 = 1.$$

So e.g. $I_2(3) = \{1, s_1, s_2, s_1 s_2, s_2 s_1, w_o = s_1 s_2 s_1 = s_2 s_1 s_2\} = S_3$.

Coxeter groups

The group S_{n+1} of permutations, and $(\mathbb{Z}/2\mathbb{Z}) \wr S_n$ of signed permutations, are examples of *finite Coxeter groups* – finite groups of orthogonal transformations generated by reflections.

- For S_{n+1} , let $s_1 = (1\ 2)$, $s_2 = (2\ 3)$, \dots , $s_n = (n\ n+1)$.

The **permutation group** S_{n+1} is generated by s_1, \dots, s_n with the **braid relations**

$$s_i s_j = s_j s_i \text{ (i.e. } (s_i s_j)^2 = 1), \quad |i - j| > 1, \quad s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1},$$

and the **Coxeter relations** $s_i^2 = 1$.

- The **dihedral group** $I_2(m)$ (rotation and reflection symmetries of the regular m -gon) is generated by s_1, s_2 with the braid and Coxeter relations

$$(s_1 s_2)^m = 1 \text{ (i.e. } s_1 s_2 s_1 \cdots = s_2 s_1 s_2 \cdots), \quad s_1^2 = s_2^2 = 1.$$

So e.g. $I_2(3) = \{1, s_1, s_2, s_1 s_2, s_2 s_1, w_o = s_1 s_2 s_1 = s_2 s_1 s_2\} = S_3$.

The finite real reflection groups were classified in 1934 by Coxeter:

Coxeter's paper

ANNALS OF MATHEMATICS
Vol. 35, No. 3, July, 1934

DISCRETE GROUPS GENERATED BY REFLECTIONS

By H. S. M. COXETER

(Received June 16, 1933)

CONTENTS

1-3. Polytopes with restricted dihedral angles.....	590
4-7. Fundamental regions.....	594
8. The abstract definition.....	599
9. Complete enumeration of the irreducible groups.....	601
10-13. Continued products of the generators.....	602
14, 15. The central inversion.....	606
16. The total number of reflections.....	610
17. Generation by two operations.....	612
18. Elliptic space.....	617
Table of irreducible finite groups generated by reflections.....	618
Graphical representation.....	619
Index.....	620
References.....	620

Introduction

It is known that, in ordinary space, the only finite groups generated by reflections are

$[k]$ ($k \geq 1$), of order $2k$, with abstract definition

$$R_1^2 = R_2^2 = (R_1 R_2)^k = 1,$$

and

$[k_1, k_2]$ ($k_1 \geq 2, k_2 \geq 2, 1/k_1 + 1/k_2 > \frac{1}{2}$), of order $\frac{4}{1/k_1 + 1/k_2 - \frac{1}{2}}$, with abstract definition

$$R_1^2 = R_2^2 = R_3^2 = (R_1 R_2)^{k_1} = (R_1 R_3)^{k_2} = (R_2 R_3)^{k_2} = 1.$$

[1] is the group of order 2 generated by a single reflection. Since this is the symmetry group of the one-dimensional polytope¹ $\{ \}$, we write

$$[1] = \{ \}.$$

$[k]$ ($k \geq 3$) and $[k_1, k_2]$ ($k_1 = 3, k_2 = 3, 4, 5$) are the symmetry groups of the ordinary regular polygons $\{k\}$ and polyhedra $\{k_1, k_2\}$. The rest of the groups can be written as direct products, thus:

$$[2] = \{ \} \times \{ \},$$

TABLE OF IRREDUCIBLE FINITE GROUPS GENERATED BY REFLECTIONS

Group	Order ²	m	k	Central inversion?
$[3^*]$	$(n+2)!$	$n+1$	$n+2$	Only when $n=0$
$[3^*, 4]$	$2^{n+2}(n+2)!$	$n+2$	$2(n+2)$	Yes
$\begin{bmatrix} 3^* \\ 3 \\ 3 \end{bmatrix}$	$2^{n+2}(n+3)!$	$n+3$	$2(n+2)$	Only when n is odd
$[k]$	$2k$	2	k	Only when k is even
$[3, 5]$	120	3	10	Yes
$[3, 4, 3]$	1152	4	12	Yes
$[3, 3, 5]$	14400	4	30	Yes
$\begin{bmatrix} 3, 3 \\ 3, 3 \\ 3 \end{bmatrix}$	51840	6	12	No
$\begin{bmatrix} 3, 3, 3 \\ 3, 3 \\ 3 \end{bmatrix}$	2903040	7	18	Yes
$\begin{bmatrix} 3, 3, 3, 3 \\ 3, 3 \\ 3 \end{bmatrix}$	696729600	8	30	Yes

Presentation of finite Coxeter group(algebra)s

Every (finite) Coxeter group W is given by:

- A finite set of generators $\{s_i \mid i \in I\}$.
- A symmetric Coxeter integer matrix $M = (m_{ij})_{i,j \in I}$, with $2 = m_{ii} \leq m_{ij} \leq \infty$.

Presentation of finite Coxeter group(algebra)s

Every (finite) Coxeter group W is given by:

- A finite set of generators $\{s_i \mid i \in I\}$.
- A symmetric Coxeter integer matrix $M = (m_{ij})_{i,j \in I}$, with $2 = m_{ii} \leq m_{ij} \leq \infty$.
- The *braid relations* $(s_i s_j)^{m_{ij}} = 1 \ \forall i \neq j$.
- The *Coxeter relations* $s_i^2 = 1 \ \forall i$.

Presentation of finite Coxeter group(algebra)s

Every (finite) Coxeter group W is given by:

- A finite set of generators $\{s_i \mid i \in I\}$.
- A symmetric Coxeter integer matrix $M = (m_{ij})_{i,j \in I}$, with $2 = m_{ii} \leq m_{ij} \leq \infty$.
- The *braid relations* $(s_i s_j)^{m_{ij}} = 1 \ \forall i \neq j$.
- The *Coxeter relations* $s_i^2 = 1 \ \forall i$.

Associated to W is its *group algebra* $\mathbb{k}W$ for any field \mathbb{k} (or even unital commutative ring). Clearly, $\dim \mathbb{k}W = |W|$.

Presentation of finite Coxeter group(algebra)s

Every (finite) Coxeter group W is given by:

- A finite set of generators $\{s_i \mid i \in I\}$.
- A symmetric Coxeter integer matrix $M = (m_{ij})_{i,j \in I}$, with $2 = m_{ii} \leq m_{ij} \leq \infty$.
- The *braid relations* $(s_i s_j)^{m_{ij}} = 1 \ \forall i \neq j$.
- The *Coxeter relations* $s_i^2 = 1 \ \forall i$.

Associated to W is its *group algebra* $\mathbb{k}W$ for any field \mathbb{k} (or even unital commutative ring). Clearly, $\dim \mathbb{k}W = |W|$.

There exist other algebras with the same dimension, which are “deformations” of $\mathbb{k}W$:

From the braid monoid-algebra...

Given a Coxeter system I and matrix $M \in \mathbb{Z}_{\geq 2}^{I \times I}$,

- The free associative algebra on I is $\mathbb{k}\langle T_i \mid i \in I \rangle$, with basis all words in the generators T_i . (Equivalently, the tensor algebra over \mathbb{k} of $\bigoplus_i (\mathbb{k}T_i)$.)
- The corresponding *monoid algebra* is its quotient by a two-sided ideal:

$$\mathbb{k}\mathcal{B}_M := \mathbb{k}\langle T_i \mid i \in I \rangle / (T_i T_j T_i \cdots = T_j T_i T_j \cdots \mid i \neq j);$$

ANNALS OF MATHEMATICS
 Vol. 48, No. 1, January, 1947

THEORY OF BRAIDS

By E. ARTIN

(Received May 20, 1946)

A theory of braids leading to a classification was given in my paper "Theorie der Zöpfe" in vol. 4 of the *Hamburger Abhandlungen* (quoted as *Z*). Most of the proofs are entirely intuitive. That of the main theorem in §7 is not even convincing. It is possible to correct the proofs. The difficulties that one encounters if one tries to do so come from the fact that projection of the braid, which is an excellent tool for intuitive investigations, is a very clumsy one for rigorous proofs. This has led me to abandon projections altogether. We shall use the more powerful tool of braid coordinates and obtain thereby farther reaching results of greater generality.

... to generic Hecke algebras

- The associated *generic Hecke algebra* (with parameters $a, b \in \mathbb{k}$) is:

$$\mathcal{E}_{a,b} := \mathbb{k}\mathcal{B}_M / (T_i^2 - aT_i - b \mid i \in I).$$

- Special case: the *Iwahori–Hecke algebras* $\mathcal{H}_q(W)$ that are prominent in representation theory; here $a = q - 1$, $b = q$.
(As $q \rightarrow 1$, say over $\mathbb{k} = \mathbb{R}$, the relations go to $\mathbb{k}W$: “deformation”.)

... to generic Hecke algebras

- The associated *generic Hecke algebra* (with parameters $a, b \in \mathbb{k}$) is:

$$\mathcal{E}_{a,b} := \mathbb{k}\mathcal{B}_M / (T_i^2 - aT_i - b \mid i \in I).$$

- Special case: the *Iwahori-Hecke algebras* $\mathcal{H}_q(W)$ that are prominent in representation theory; here $a = q - 1$, $b = q$.
(As $q \rightarrow 1$, say over $\mathbb{k} = \mathbb{R}$, the relations go to $\mathbb{k}W$: “deformation”.)

Fact: Each such algebra $\mathcal{E}_{a,b}$ has a “word basis” $\{T_w : w \in W\}$.
So, its dimension is $|W|$.

Nil-Coxeter algebras

There are three special cases of (a, b) which are interesting from the viewpoint of combinatorics, PBW deformation theory, ...:

- ❶ $a = 0, b = 1$ – group algebra $\mathbb{k}W$.
- ❷ $a = 1, b = 0$ – 0-Hecke algebra (Norton, Hivert–Schilling–Thiery, ...).
- ❸ $a = b = 0$ – **nil-Coxeter algebra** NC_W . So, $T_i^2 = 0$ – *graded* algebra.

Nil-Coxeter algebras

There are three special cases of (a, b) which are interesting from the viewpoint of combinatorics, PBW deformation theory, ...:

- ① $a = 0, b = 1$ – group algebra $\mathbb{k}W$.
- ② $a = 1, b = 0$ – 0-Hecke algebra (Norton, Hivert–Schilling–Thiery, ...).
- ③ $a = b = 0$ – **nil-Coxeter algebra** NC_W . So, $T_i^2 = 0$ – *graded* algebra.

Nil-Coxeter algebras “occur naturally” as differential / divided-difference operators on polynomial rings (and hence in Schubert calculus). E.g. in type A , for $1 \leq i \leq n$ the operator T_i is:

$$(T_i f)(x_1, \dots, x_{n+1}) = \frac{f(x_1, \dots, x_i, x_{i+1}, \dots, x_{n+1}) - f(x_1, \dots, x_{i+1}, x_i, \dots, x_{n+1})}{x_{i+1} - x_i}.$$

(The RHS is $\frac{\text{alternating}}{\text{alternating}}$ = symmetric in $\{x_i, x_{i+1}\}$, so $T_i^2 f = 0$.)

Word basis of NC_W

Example: The dihedral group D_n (e.g. S_3) has elements

$$e, s_1, s_2, s_1s_2, s_2s_1, s_1s_2s_1, s_2s_1s_2, \dots,$$

increasing in length all the way to the unique longest element

$$s_1s_2s_1 \cdots = w_o = s_2s_1s_2 \cdots .$$

Correspondingly,

Word basis of NC_W

Example: The dihedral group D_n (e.g. S_3) has elements

$$e, s_1, s_2, s_1s_2, s_2s_1, s_1s_2s_1, s_2s_1s_2, \dots,$$

increasing in length all the way to the unique longest element

$$s_1s_2s_1 \cdots = w_o = s_2s_1s_2 \cdots .$$

Correspondingly, its nil-Coxeter algebra has a “word basis”

$$1 = T_\emptyset, T_1, T_2, T_1T_2, T_2T_1, T_1T_2T_1, T_2T_1T_2, \dots,$$

all the way to the unique longest element T_{w_o} .

Now come to the protagonist of this talk: “generalized” nil-Coxeter algebras.

Generalized Coxeter groups

In 1957, Coxeter studied the “generalized” Coxeter group $W_A(n, d)$, defined as the quotient of the type- A (Artin) braid monoid by $s_i^d = 1 \ \forall i$.
(Such higher orders are typical in *complex* reflection groups.)

- Coxeter asked: *For which $n, d \geq 2$ is $W_A(n, d)$ a finite group?*

Generalized Coxeter groups

In 1957, Coxeter studied the “generalized” Coxeter group $W_A(n, d)$, defined as the quotient of the type- A (Artin) braid monoid by $s_i^d = 1 \ \forall i$.
(Such higher orders are typical in *complex* reflection groups.)

- Coxeter asked: *For which $n, d \geq 2$ is $W_A(n, d)$ a finite group?*

Theorem (Coxeter, 1957)

$W_A(n, d)$ is a finite group if and only if $\frac{1}{n} + \frac{1}{d} > \frac{1}{2}$, and then
 $|W_A(n, d)| = \left(\frac{1}{n} + \frac{1}{d} - \frac{1}{2}\right)^{1-n} n! / n^{n-1}$.

Generalized Coxeter groups

In 1957, Coxeter studied the “generalized” Coxeter group $W_A(n, d)$, defined as the quotient of the type- A (Artin) braid monoid by $s_i^d = 1 \ \forall i$.
 (Such higher orders are typical in *complex* reflection groups.)

- Coxeter asked: *For which $n, d \geq 2$ is $W_A(n, d)$ a finite group?*

Theorem (Coxeter, 1957)

$W_A(n, d)$ is a finite group if and only if $\frac{1}{n} + \frac{1}{d} > \frac{1}{2}$, and then
 $|W_A(n, d)| = \left(\frac{1}{n} + \frac{1}{d} - \frac{1}{2}\right)^{1-n} n! / n^{n-1}$.

Later extended by Koster to cover all generalized Coxeter groups.

Generalized nil-Coxeter algebras

Note: In Coxeter's construction, all s_i are conjugate because of the braid relations. Hence their orders are equal.

Generalized nil-Coxeter algebras

Note: In Coxeter's construction, all s_i are conjugate because of the braid relations. Hence their orders are equal.

Not necessary in the nil-Coxeter case. Thus:

- Given integers $d_i \geq 2$, **define** the *generalized nil-Coxeter algebra* (over W or M) to be

$$NC_M(\{d_i \mid i \in I\}) := \mathbb{k}\mathcal{B}_M / (T_i^{d_i} = 0 \mid i \in I).$$

(Still a $\mathbb{Z}_{\geq 0}$ -graded algebra.)

Generalized nil-Coxeter algebras

Note: In Coxeter's construction, all s_i are conjugate because of the braid relations. Hence their orders are equal.

Not necessary in the nil-Coxeter case. Thus:

- Given integers $d_i \geq 2$, **define** the *generalized nil-Coxeter algebra* (over W or M) to be

$$NC_M(\{d_i \mid i \in I\}) := \mathbb{k}\mathcal{B}_M / (T_i^{d_i} = 0 \mid i \in I).$$

(Still a $\mathbb{Z}_{\geq 0}$ -graded algebra.)

- Question** (*a la Coxeter*): For which Coxeter groups $W = (I, M)$ and tuples $\mathbf{d} = (d_i)_i$ is this algebra $NC_M(\mathbf{d})$ finite-dimensional? Does it have a word basis?

Generalized nil-Coxeter algebras

Note: In Coxeter's construction, all s_i are conjugate because of the braid relations. Hence their orders are equal.

Not necessary in the nil-Coxeter case. Thus:

- Given integers $d_i \geq 2$, **define** the *generalized nil-Coxeter algebra* (over W or M) to be

$$NC_M(\{d_i \mid i \in I\}) := \mathbb{k}\mathcal{B}_M / (T_i^{d_i} = 0 \mid i \in I).$$

(Still a $\mathbb{Z}_{\geq 0}$ -graded algebra.)

- Question** (*a la Coxeter*): For which Coxeter groups $W = (I, M)$ and tuples $\mathbf{d} = (d_i)_i$ is this algebra $NC_M(\mathbf{d})$ finite-dimensional? Does it have a word basis?
- Clearly, the “usual” nil-Coxeter algebras $NC_M((2, \dots, 2))$ work.
 Any others?

Additional motivations / properties of generalized nil-Coxeter algebras

- ① *Coxeter combinatorics*: Parallel to Coxeter's question.
- ② *Tensor categories*: Generalized nil-Coxeter algebras $NC_M(\mathbf{d})$ possess a coproduct $\Delta(T_i) = T_i \otimes T_i$.
 - This is an algebra map, but cannot have a counit or antipode.
 - Thus, these are not bialgebras; hence their modules do not form a monoidal / tensor category.
 - Yet, the Tannakian formalism applies – to yield *semigroup categories* – no unit object.¹

¹This is a semigroup category under \otimes , which is additive and with \otimes bi-additive.

Additional motivations / properties of generalized nil-Coxeter algebras

- ❶ *Coxeter combinatorics*: Parallel to Coxeter's question.
- ❷ *Tensor categories*: Generalized nil-Coxeter algebras $NC_M(\mathbf{d})$ possess a coproduct $\Delta(T_i) = T_i \otimes T_i$.
 - This is an algebra map, but cannot have a counit or antipode.
 - Thus, these are not bialgebras; hence their modules do not form a monoidal / tensor category.
 - Yet, the Tannakian formalism applies – to yield *semigroup categories* – no unit object.¹
- ❸ *PBW deformations*: Despite no counit or antipode, generalized nil-Coxeter algebras $NC_M(\mathbf{d})$ smash-product polynomial rings admit “PBW deformations”.
 (Going beyond the “traditional” PBW program in the literature – Etingof–Ginzburg, Shepler–Witherspoon (and Walton), ...)

¹This is a semigroup category under \otimes , which is additive and with \otimes bi-additive.

A novel type- A family

Back to the question: Classify the finite-dimensional generalized nil-Coxeter algebras $NC_M(\mathbf{d})$ for $\mathbf{d} \in \mathbb{Z}_{\geq 2}^I$ – outside of the “usual” nil-Coxeter algebras with all $d_i = 2$.

A novel type- A family


Back to the question: Classify the finite-dimensional generalized nil-Coxeter algebras $NC_M(\mathbf{d})$ for $\mathbf{d} \in \mathbb{Z}_{\geq 2}^I$ – outside of the “usual” nil-Coxeter algebras with all $d_i = 2$.

- An obvious positive answer is: $NC_A(1, d) := \mathbb{k}[T_1]/(T_1^d)$.

A novel type- A family

Back to the question: Classify the finite-dimensional generalized nil-Coxeter algebras $NC_M(\mathbf{d})$ for $\mathbf{d} \in \mathbb{Z}_{\geq 2}^I$ – outside of the “usual” nil-Coxeter algebras with all $d_i = 2$.

- An obvious positive answer is: $NC_A(1, d) := \mathbb{k}[T_1]/(T_1^d)$.
- This can be extended to $n = 2$: $NC_A(2, d) := \mathbb{k}\mathcal{B}_{A_2}/(T_1^2, T_2^d)$.



(In the figure, $d' = d - 1$.)

The general type- A algebra

Theorem (K., *Trans. AMS* 2018 + *FPSAC* 2018)

For every $n \geq 1$ and $d \geq 2$, the type- A generalized nil-Coxeter algebra

$$NC_A(n, d) := \mathbb{k}\mathcal{B}_{A_n} / (T_1^2, \dots, T_{n-1}^2, T_n^d)$$

is finite-dimensional (or free of finite \mathbb{k} -rank).

The general type- A algebra

Theorem (K., *Trans. AMS* 2018 + *FPSAC* 2018)

For every $n \geq 1$ and $d \geq 2$, the type- A generalized nil-Coxeter algebra

$$NC_A(n, d) := \mathbb{k}\mathcal{B}_{A_n} / (T_1^2, \dots, T_{n-1}^2, T_n^d)$$

is finite-dimensional (or free of finite \mathbb{k} -rank).

The proof involves explicitly defining an action of $NC_A(n, d)$ on its “regular representation”.

The general type- A algebra

Theorem (K., *Trans. AMS* 2018 + *FPSAC* 2018)

For every $n \geq 1$ and $d \geq 2$, the type- A generalized nil-Coxeter algebra

$$NC_A(n, d) := \mathbb{k}\mathcal{B}_{A_n} / (T_1^2, \dots, T_{n-1}^2, T_n^d)$$

is finite-dimensional (or free of finite \mathbb{k} -rank).

The proof involves explicitly defining an action of $NC_A(n, d)$ on its “regular representation”. Moreover, we compute its

- \mathbb{k} -rank: $n!(1 + n(d - 1))$.
- word basis:
 $\{T_w; T_w T_n^k T_{n-1} \cdots T_m \mid w \in S_n = W_{A_{n-1}}, k \in [d - 1], m \in [n]\}$.
- unique longest word, left/right primitive words, ...

The general type- A algebra

Theorem (K., *Trans. AMS* 2018 + *FPSAC* 2018)

For every $n \geq 1$ and $d \geq 2$, the type- A generalized nil-Coxeter algebra

$$NC_A(n, d) := \mathbb{k}\mathcal{B}_{A_n} / (T_1^2, \dots, T_{n-1}^2, T_n^d)$$

is finite-dimensional (or free of finite \mathbb{k} -rank).

The proof involves explicitly defining an action of $NC_A(n, d)$ on its “regular representation”. Moreover, we compute its

- \mathbb{k} -rank: $n!(1 + n(d - 1))$.
- word basis:
 $\{T_w; T_w T_n^k T_{n-1} \cdots T_m \mid w \in S_n = W_{A_{n-1}}, k \in [d - 1], m \in [n]\}$.
- unique longest word, left/right primitive words, ...
- The “usual” length function ℓ extends to $NC_A(n, d)$, and its Hilbert–Poincaré series (in q) is
 $[n]_q!(1 + [n]_q[d - 1]_q), \quad \text{where } [n]_q := \frac{q^n - 1}{q - 1}, [n]_q! := \prod_{j=1}^n [j]_q.$

All finite-dimensional generalized nil-Coxeter algebras

Are there any other finite-dimensional generalized nil-Coxeter algebras?

All finite-dimensional generalized nil-Coxeter algebras

Are there any other finite-dimensional generalized nil-Coxeter algebras? **No!**

Theorem (K., *Trans. AMS* 2018 + *FPSAC* 2018)

Given a Coxeter matrix $M \in \mathbb{Z}_{\geq 2}^{I \times I}$ and an integer tuple $\mathbf{d} \in \mathbb{Z}_{\geq 2}^I$, the following are equivalent:

- 1 The algebra $NC_M(\mathbf{d})$ is finite-dimensional (or of finite \mathbb{k} rank).
- 2 Either $W = W(M)$ is a finite Coxeter group and all $d_i = 2$, or W is of type A_n and $\mathbf{d} = (2, \dots, 2, d)$ (or $(d, 2, \dots, 2)$) – i.e., $NC_M(\mathbf{d}) = NC_A(n, d)$.

All finite-dimensional generalized nil-Coxeter algebras


Are there any other finite-dimensional generalized nil-Coxeter algebras? **No!**

Theorem (K., *Trans. AMS* 2018 + *FPSAC* 2018)

Given a Coxeter matrix $M \in \mathbb{Z}_{\geq 2}^{I \times I}$ and an integer tuple $\mathbf{d} \in \mathbb{Z}_{\geq 2}^I$, the following are equivalent:

- ❶ The algebra $NC_M(\mathbf{d})$ is finite-dimensional (or of finite \mathbb{k} rank).
- ❷ Either $W = W(M)$ is a finite Coxeter group and all $d_i = 2$,
 or W is of type A_n and $\mathbf{d} = (2, \dots, 2, d)$ (or $(d, 2, \dots, 2)$) – i.e.,
 $NC_M(\mathbf{d}) = NC_A(n, d)$.

The proof uses a
 diagrammatic calculus:



Complex reflection groups and the BMR Freeness Conjecture

The higher nilpotency $T_i^{d_i} = 0$ is reminiscent of *complex* reflection groups $W_{\mathbb{C}}$.

- These groups also have “Coxeter-like” presentations using nodes and edges / generators and relations. The finite groups $W_{\mathbb{C}}$ were classified by Shephard–Todd [*Canadian J. Math.* 1954].
- So, they also have *generic Hecke algebras* $\mathcal{H}_q(W_{\mathbb{C}}) \dots$

Complex reflection groups and the BMR Freeness Conjecture

The higher nilpotency $T_i^{d_i} = 0$ is reminiscent of *complex* reflection groups $W_{\mathbb{C}}$.

- These groups also have “Coxeter-like” presentations using nodes and edges / generators and relations. The finite groups $W_{\mathbb{C}}$ were classified by Shephard–Todd [*Canadian J. Math.* 1954].
- So, they also have *generic Hecke algebras* $\mathcal{H}_q(W_{\mathbb{C}}) \dots$
- ... akin to which, one forms generalized nil-Coxeter algebras $NC_{W_{\mathbb{C}}}$.

Which of these algebras $\mathcal{H}_q(W_{\mathbb{C}})$ and $NC_{W_{\mathbb{C}}}$ are finite-dimensional? Of dimension $W_{\mathbb{C}}$?

Complex reflection groups and the BMR Freeness Conjecture

The higher nilpotency $T_i^{d_i} = 0$ is reminiscent of *complex* reflection groups $W_{\mathbb{C}}$.

- These groups also have “Coxeter-like” presentations using nodes and edges / generators and relations. The finite groups $W_{\mathbb{C}}$ were classified by Shephard–Todd [*Canadian J. Math.* 1954].
- So, they also have *generic Hecke algebras* $\mathcal{H}_q(W_{\mathbb{C}}) \dots$
- ... akin to which, one forms generalized nil-Coxeter algebras $NC_{W_{\mathbb{C}}}$.

Which of these algebras $\mathcal{H}_q(W_{\mathbb{C}})$ and $NC_{W_{\mathbb{C}}}$ are finite-dimensional? Of dimension $W_{\mathbb{C}}$?

Broué–Malle–Rouquier Freeness Conjecture (Crelle 1998)

Generic Hecke algebras \mathcal{H}_q over $(W_{\mathbb{C}}, \mathbb{k})$ are free with \mathbb{k} -rank $|W_{\mathbb{C}}|$.

(Proved by Etingof in 2017, in characteristic zero.)

Generalized nil-Coxeter algebras over $W_{\mathbb{C}}$

What about (generalized) nil-Coxeter algebras?

Generalized nil-Coxeter algebras over $W_{\mathbb{C}}$

What about (generalized) nil-Coxeter algebras?

Already the nil-Coxeter picture was “discouraging”: Ivan Marin had written [JPAA 2014] that “the lack of nil-Coxeter algebras of dimension $|W_{\mathbb{C}}|$ is a striking difference between real and complex reflection groups.”

Generalized nil-Coxeter algebras over $W_{\mathbb{C}}$

What about (generalized) nil-Coxeter algebras?

Already the nil-Coxeter picture was “discouraging”: Ivan Marin had written [JPAA 2014] that “the lack of nil-Coxeter algebras of dimension $|W_{\mathbb{C}}|$ is a striking difference between real and complex reflection groups.”

This was actually **Motivation 4** for us. And we showed (2018):

- *There are no finite-dimensional (let alone $|W_{\mathbb{C}}|$ -dim.) **generalized** nil-Coxeter algebras over $W_{\mathbb{C}}$.*

Generalized nil-Coxeter algebras over $W_{\mathbb{C}}$

What about (generalized) nil-Coxeter algebras?

Already the nil-Coxeter picture was “discouraging”: Ivan Marin had written [JPAA 2014] that “the lack of nil-Coxeter algebras of dimension $|W_{\mathbb{C}}|$ is a striking difference between real and complex reflection groups.”

This was actually **Motivation 4** for us. And we showed (2018):

- *There are no finite-dimensional (let alone $|W_{\mathbb{C}}|$ -dim.) **generalized** nil-Coxeter algebras over $W_{\mathbb{C}}$.*

Upshot: The novel family $NC_A(n, d)$ is strikingly unique, among all real and complex reflection groups!

Does it “occur in nature” (akin to the divided difference operators for $d = 2$)?

Nil-Temperley–Lieb algebras

What next?

Nil-Temperley–Lieb algebras

What next?

Kill long enough braid words!

The *Temperley–Lieb algebra*
in types A, D, E is
defined as the quotient of the
Iwahori–Hecke algebra by the
ideal generated by

$T_s T_t T_s = \text{lower degree terms}$

for adjacent nodes s, t
in the Coxeter graph.

JOURNAL OF THE
AMERICAN MATHEMATICAL SOCIETY
Volume 10, Number 1, January 1997, Pages 139–167
S 0894-0347(97)00222-1

STRUCTURE OF A HECKE ALGEBRA QUOTIENT

C. KENNETH FAN

Dedicated to my teacher, George Lusztig, on his fiftieth birthday

1. INTRODUCTION

Let W be a Coxeter group with Coxeter graph Γ . Let Γ_g be the set of simple generators, which are parametrized by the nodes of Γ .

Our primary interest in this paper is to understand the case where Γ is of type E . Therefore, we shall assume that Γ is of type A, D , or E , where by E , we mean the infinite series E_n which begins $E_5 = D_5, E_6, E_7, E_8, E_9 = \hat{E}_8$, etc.

Every $w \in W$ may be written as a product $s_1 s_2 s_3 \cdots s_n$ of generators in Γ_g . If n is minimal, we call this product “reduced” and define $l(w) = n$. More generally, if $w = w_1 w_2 w_3 \cdots w_n$ satisfies $l(w) = \sum_i l(w_i)$, then we call this product “reduced” as well.

Let \mathcal{H} be the Iwahori–Hecke algebra associated to W . This is an algebra over $\mathbb{Q}(q^{1/2})$ (where $q^{1/2}$ is an indeterminate) with generators T_s for each $s \in \Gamma_g$ satisfying the relations $T_s^2 = (q-1)T_s + q$, $T_s T_t = T_t T_s$ if $st = ts$, and $T_s T_t T_s = T_t T_s T_t$ if $sts = tst$, where $s, t \in \Gamma_g$. This algebra has a basis T_w , $w \in W$, where we have $T_w = T_{s_1} \cdots T_{s_n}$ whenever $s_1 \cdots s_n$ is a reduced expression for w .

Let \mathcal{I} be the two-sided ideal generated by the elements

$$T_{sts} + T_{st} + T_{ts} + T_s + T_t + 1$$

Nil-Temperley–Lieb algebras

There is a nil-version: given a Coxeter group W with data (I, M) , the *nil-Temperley–Lieb algebra* $NTL_W = NTL_M$ is the quotient of $\mathbb{k}\mathcal{B}_M$ by

- the “braid relations” $T_s T_t T_s = 0$ for adjacent nodes $s \sim t$;
- and the Coxeter relations $T_i^2 = 0$ for all $i \in I$.

Nil-Temperley–Lieb algebras

There is a nil-version: given a Coxeter group W with data (I, M) , the *nil-Temperley–Lieb algebra* $NTL_W = NTL_M$ is the quotient of $\mathbb{k}\mathcal{B}_M$ by

- the “braid relations” $T_s T_t T_s = 0$ for adjacent nodes $s \sim t$;
- and the Coxeter relations $T_i^2 = 0$ for all $i \in I$.

These were (re?)introduced by A. Postnikov as *XYX-algebras*.

Clearly, “usual” nil-Coxeter algebras surject onto them.

Which of them are finite-dimensional?

Nil-Temperley–Lieb algebras

There is a nil-version: given a Coxeter group W with data (I, M) , the *nil-Temperley–Lieb algebra* $NTL_W = NTL_M$ is the quotient of $\mathbb{k}\mathcal{B}_M$ by

- the “braid relations” $T_s T_t T_s = 0$ for adjacent nodes $s \sim t$;
- and the Coxeter relations $T_i^2 = 0$ for all $i \in I$.


These were (re?)introduced by A. Postnikov as *XYX-algebras*.

Clearly, “usual” nil-Coxeter algebras surject onto them.

Which of them are finite-dimensional?

Theorem (essentially due to Stembridge, C.K. Fan, 1990s)

NTL_W has finite \mathbb{k} -rank if and only if W is a finite Coxeter group, or W has one of the following Coxeter graphs:



(Generalized) nil-Temperley–Lieb algebras

Question: In the *generalized* nil-Temperley–Lieb version, with relations $T_s T_t T_s = 0$ and $T_i^{d_i} = 0$, which algebras $NTL_M(\mathbf{d})$ are finite-dimensional?

(Generalized) nil-Temperley–Lieb algebras

Question: In the *generalized* nil-Temperley–Lieb version, with relations $T_s T_t T_s = 0$ and $T_i^{d_i} = 0$, which algebras $NTL_M(\mathbf{d})$ are finite-dimensional?

Theorem (Bhattacharyya–K., 2021 preprint)

If and only if

- 1 The algebras on the previous slide;
- 2 “Generalized XYX -algebras”
 $NTL_A(n, d) := NC_A(n, d) / (T_s T_t T_s, |s - t| = 1).$

(Generalized) nil-Temperley–Lieb algebras

Question: In the *generalized* nil-Temperley–Lieb version, with relations $T_s T_t T_s = 0$ and $T_i^{d_i} = 0$, which algebras $NTL_M(\mathbf{d})$ are finite-dimensional?

Theorem (Bhattacharyya–K., 2021 preprint)

If and only if

- 1 The algebras on the previous slide;
- 2 “Generalized XYX -algebras”
 $NTL_A(n, d) := NC_A(n, d) / (T_s T_t T_s, |s - t| = 1).$

In the first case, the dimension is $|W_{fc}| < \infty$, the fully commutative words.^a

(Generalized) nil-Temperley–Lieb algebras

Question: In the *generalized* nil-Temperley–Lieb version, with relations $T_s T_t T_s = 0$ and $T_i^{d_i} = 0$, which algebras $NTL_M(\mathbf{d})$ are finite-dimensional?

Theorem (Bhattacharyya–K., 2021 preprint)

If and only if

- 1 The algebras on the previous slide;
- 2 “Generalized XYX -algebras”

$$NTL_A(n, d) := NC_A(n, d) / (T_s T_t T_s, |s - t| = 1).$$

In the first case, the dimension is $|W_{fc}| < \infty$, the fully commutative words.^a

In the second case,

$$\dim NTL_A(n, d) = (d - 1)C_{n+1} - (d - 2)C_n + (d - 2) \sum_{j=1}^{n-1} jC_{n-j}, \quad (1.6)$$

where C_n is the n th Catalan number.

^aThe words in W for which switching between any two reduced expressions uses no non-commutative braid relations.

Going beyond type A

Similarly, one can quotient by all braid words of **length ≥ 4** (but not the braid words $T_s T_t T_s$):

- In the simply-laced types A, D, E , this simply yields the “usual” nil-Coxeter algebras (since no “extra” quotienting is needed).
- Ditto in the case of the algebras $NC_A(n, d)$ – they remain untouched.

Going beyond type A

Similarly, one can quotient by all braid words of **length ≥ 4** (but not the braid words $T_s T_t T_s$):

- In the simply-laced types A, D, E , this simply yields the “usual” nil-Coxeter algebras (since no “extra” quotienting is needed).
- Ditto in the case of the algebras $NC_A(n, d)$ – they remain untouched.
- In type B_n , the words in the T_i that remain nonzero precisely correspond to the $\overline{1} \overline{2}$ -avoiding signed permutations – there are $\sum_{k=0}^n \binom{n}{k}^2 k!$ many.

Going beyond type A

Similarly, one can quotient by all braid words of **length ≥ 4** (but not the braid words $T_s T_t T_s$):

- In the simply-laced types A, D, E , this simply yields the “usual” nil-Coxeter algebras (since no “extra” quotienting is needed).
- Ditto in the case of the algebras $NC_A(n, d)$ – they remain untouched.
- In type B_n , the words in the T_i that remain nonzero precisely correspond to the $\overline{1} \overline{2}$ -avoiding signed permutations – there are $\sum_{k=0}^n \binom{n}{k}^2 k!$ many.

Similar results if one quotients by the braid words of **length ≥ 5** .

- There is exactly one missing case: H_n for $n \geq 5$ (equivalently in the **length ≥ 4** case, because there is only one such pair of words: $T_1 T_2 T_1 T_2 T_1 = T_2 T_1 T_2 T_1 T_2$ – see the Figure on Slide 18).

Going beyond type A

Similarly, one can quotient by all braid words of **length ≥ 4** (but not the braid words $T_s T_t T_s$):

- In the simply-laced types A, D, E , this simply yields the “usual” nil-Coxeter algebras (since no “extra” quotienting is needed).
- Ditto in the case of the algebras $NC_A(n, d)$ – they remain untouched.
- In type B_n , the words in the T_i that remain nonzero precisely correspond to the $\bar{1}\bar{2}$ -avoiding signed permutations – there are $\sum_{k=0}^n \binom{n}{k}^2 k!$ many.

Similar results if one quotients by the braid words of **length ≥ 5** .

- There is exactly one missing case: H_n for $n \geq 5$ (equivalently in the **length ≥ 4** case, because there is only one such pair of words: $T_1 T_2 T_1 T_2 T_1 = T_2 T_1 T_2 T_1 T_2$ – see the Figure on Slide 18).

If instead one kills all braid words of **length ≥ 6** , then since there were no such in any non-dihedral finite Coxeter group, we just get back the usual nil-Coxeter algebra (or the algebras $NC_A(n, d)$) – hence of finite rank.

Table of findings (from the preprint)

Table of all finite-dimensional generalized nil-Temperley–Lieb algebras.

In it, $J_{<k}$ means we quotient by all braid words of length $\geq k$.

W	$J_{<3}$	$J_{<4}$	$J_{<5}$	$J_{<k} \ (6 \leq k \leq \infty)$
A_n, D_n, E_6, E_7, E_8	$ W_{\text{fc}} $	$ W $	$ W $	$ W $
B_n	$ W_{\text{fc}} $	$\sum_{k=0}^n \binom{n}{k}^2 k!$	$ W $	$ W $
F_4	$ W_{\text{fc}} $	304	$ W $	$ W $
H_3	$ W_{\text{fc}} $	76	76	$ W $
H_4	$ W_{\text{fc}} $	1460	1460	$ W $
$I_2(m)$	$ W $ if $m < 3$, else $ W_{\text{fc}} = W - 1$	$ W $ if $m < 4$, else $ W - 1$	$ W $ if $m < 5$, else $ W - 1$	$ W $ if $m < k$, else $ W - 1$
$E_n \ (n \geq 9)$	$ W_{\text{fc}} $	–	–	–
$F_n \ (n \geq 5)$	$ W_{\text{fc}} $	–	–	–
$H_n \ (n \geq 5)$	$ W_{\text{fc}} $?	?	–
$A_n, \mathbf{d} = (d, 2, \dots, 2),$ $d > 2$	(see (1.6))	$n!(1 + n(d - 1))$	$n!(1 + n(d - 1))$	$n!(1 + n(d - 1))$

Open questions

- 1 Do the algebras $NC_A(n, d)$ “occur in nature”? How about the generalized XYX -algebras $NTL_A(n, d)$ with $d \geq 3$?

Open questions

- 1 Do the algebras $NC_A(n, d)$ “occur in nature”? How about the generalized XYX -algebras $NTL_A(n, d)$ with $d \geq 3$?
- 2 Find a combinatorial way to enumerate the word basis of $NTL_A(n, d)$; recall this has size

$$\dim NTL_A(n, d) = (d-1)C_{n+1} - (d-2)C_n + (d-2) \sum_{j=1}^{n-1} jC_{n-j}.$$

Open questions

- 1 Do the algebras $NC_A(n, d)$ “occur in nature”? How about the generalized XYX -algebras $NTL_A(n, d)$ with $d \geq 3$?

- 2 Find a combinatorial way to enumerate the word basis of $NTL_A(n, d)$; recall this has size

$$\dim NTL_A(n, d) = (d-1)C_{n+1} - (d-2)C_n + (d-2) \sum_{j=1}^{n-1} jC_{n-j}.$$

- 3 For which $n \geq 5$ does the H_n nil-Temperley–Lieb algebra become of finite \mathbb{k} -rank, when one quotients by braid words of length ≥ 4 (equivalently, length ≥ 5 – that is, $T_1T_2T_1T_2T_1$ and $T_2T_1T_2T_1T_2$) ?

References

- [1] A. Khare.
Generalized nil-Coxeter algebras over discrete complex reflection groups.
Trans. Amer. Math. Soc., 2018 (+ FPSAC 2018).

- [2] Sutanay Bhattacharya and A. Khare.
The lattice of nil-Hecke algebras over real and complex reflection groups.
Preprint (under revision), 2021.