

# 1 Recap

1. Determinant of exponentials.
2. Solution of homogeneous and inhomogeneous autonomous problems.
3. Uniqueness for non-autonomous ones.

## 2 Linear systems

For the inhomogeneous non-autonomous case,  $y' = A(t)y + B(t)$ , Suppose we solve the homogenous system  $y' = A(t)y$  with  $y(t_0) = e_i(t_0)$  (note that we already know uniqueness, and we are assuming existence), we get a bunch of solutions that are linearly independent (why?) and if arrange them in a column, we get an invertible matrix  $\Phi(t, t_0)$ . Let  $\Psi(t)$  be any invertible matrix satisfying  $\Psi' = A(t)\Psi$ . Note that  $\Phi(t, t_0) = \Psi(t)\Psi(t_0)^{-1}$  (why?) As a consequence,  $\vec{y}(t) = \Phi(t, t_0)\vec{y}(t_0)$  and hence the space of solutions is  $n$ -dimensional.

We claim that the solution to  $y' = A(t)y + B(t)$  is  $y(t) = \Phi(t, t_0)y_0 + \int_{t_0}^t \Phi(t, s)B(s)ds$  akin to the Duhamel formula for the autonomous case.

*Proof.* Let's differentiate this formula and check that it satisfies the equation (it obviously satisfies the initial conditions).

$$\begin{aligned} y' &= A\Phi(t, t_0)y_0 + \frac{d}{dt} \int_{t_0}^t \Phi(t, s)B(s)ds = Ay + \Psi' \int_{t_0}^t \Psi(s)^{-1}B(s)ds \\ &= A\Phi(t, t_0)y_0 + A\Psi(t) \int_{t_0}^t \Psi^{-1}(s)B(s)ds = Ay. \end{aligned} \tag{1}$$

□

## 3 Real-analytic functions

The method of exponentiation tells us that it might be prudent to try to solve ODE using power series. To this end, we make a definition:

Def: A function  $f : (a, b) \rightarrow \mathbb{R}$  is said to be real-analytic at  $t_0$  if there exists a  $\delta > 0$  such that  $f(t) = \sum_{k=0}^{\infty} c_k(t - t_0)^k$  converges for all  $t \in (t_0 - \delta, t_0 + \delta)$ . It is said to be real-analytic on  $(a, b)$  if it is real-analytic at every point in  $(a, b)$ .

Before we come up with examples, here are some important facts from analysis:

1. The Ratio test: Let  $L = \limsup \frac{|a_{n+1}|}{|a_n|}$  and  $l$  be the lim inf. If  $L < 1$  then  $\sum a_n$  converges absolutely. If  $l > 1$ , it diverges.
2. The Root test: Let  $L = \limsup |a_n|^{1/n}$ . If  $L < 1$  then  $\sum a_n$  converges absolutely. If  $L > 1$  it diverges.

3. Applying the root test to power series we see that if  $R^{-1} = \limsup |a_n|^{1/n} > 0$ , then  $\sum a_n x^n$  converges absolutely when  $|x| < R$  (and uniformly on any compact subset of the disc of convergence) and diverges when  $|x| > R$ . This  $R$  is called the radius of convergence. (The ratio test can also be used to determine  $R$  in many cases.)
4. On a compact subset of the disc of convergence, the power series can be differentiated and integrated term-by-term to get a power series that also converges uniformly.