

NOTES FOR 9 SEPT (TUESDAY)

1. RECAP

- (1) Defined induced connections and metric compatible ones.
- (2) Defined PDE.
- (3) Levi-Civita connection existence and uniqueness. Stated theorem on the torsion tensor.

2. LEVI-CIVITA CONNECTION

Theorem 2.1. *Suppose M is a manifold. Let ∇^* be the induced connection on T^*M from any connection on TM . Then $d^{\nabla^*} : \Omega^1(M) \rightarrow \Omega^2(M)$.*

- (1) $(d^{\nabla^*} - d)\omega$ satisfies tensoriality and hence there exists a tensor $T \in \Gamma(T^{**}M \simeq TM \otimes \Omega^2(M))$ such that $T_\omega(-, -) = (d^{\nabla^*} - d)(\omega)$.
- (2) $T(X, Y) = \omega(\nabla_X Y - \nabla_Y X - [X, Y])$. Thus for the Levi-Civita connection, $d^{\nabla^*} = d$.

Proof. (1) Already done.

(2) Suppose

$$\begin{aligned}
 T_\omega(X, Y) &= T_\omega(X^i \frac{\partial}{\partial x^i}, Y^j \frac{\partial}{\partial x^j}) = X^i Y^j T(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}) = X^i Y^j (d^{\nabla^*} - d)(\omega)(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}) \\
 &= X^i Y^j (d^{\nabla^*} - d)(\omega_k dx^k)(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}) = X^i Y^j (A^*)_k^a \wedge \omega_a dx^k ((\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j})) \\
 &= X^i Y^j (\delta_j^k (A^*)_k^a (\frac{\partial}{\partial x^i}) \omega_a - \delta_i^k (A^*)_k^a (\frac{\partial}{\partial x^j}) \omega_a) = X^i Y^j (\omega_a \left(-A_j^a (\frac{\partial}{\partial x^i}) + A_i^a (\frac{\partial}{\partial x^j}) \right)) \\
 (2.1) \qquad &= \omega(\nabla_X Y - \nabla_Y X - [X, Y])
 \end{aligned}$$

□

As for the connection and curvature matrices for the Levi-Civita connection, $A_{-j}^i = \Gamma_{j\mu}^i dx^\mu$ and $F_j^i = F_{j\mu\nu}^i dx^\mu \wedge dx^\nu$ which is a complicated expression involving two derivatives of the metric g . Typically, one writes R (standing for Riemann) for the curvature tensor instead of F . It satisfies various symmetries. It also satisfies the following expression (which is usually given as a definition).

$$(2.2) \qquad R(X, Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z.$$

So the curvature matrix is $F_{-j}^i = dx^i (R(\cdot, \cdot) \frac{\partial}{\partial x^j})$. While the curvature is important, it is too much information to keep track of. Here are other “curvatures” derived from this basic object that may sometimes be easier to handle.

- (1) Sectional curvature of a two-plane spanned by an orthonormal set $X, Y \in T_p M$: This quantity is $g(R(X, Y)Y, X)$. It turns out that it is independent of the orthonormal frame chosen (using the symmetries of the Riemann tensor) and actually, using a polarisation-type identity, one can know the full Riemann tensor if one knows all sectional curvatures. Using this concept one can talk of “positively curved” (all sectional curvatures are positive everywhere) or “constant curvature” (all sectional curvatures are constant). For instance, we have a few important theorems.

- (a) Complete Riemannian manifolds with constant sectional curvature are an isometric quotient of space forms : Hyperbolic space, or Euclidean space, or the Sphere. (Killing-Hopf theorem)
 - (b) If the sectional curvature of a complete manifold is non-positive, then the universal cover is diffeomorphic to \mathbb{R}^n (Cartan-Hadamard theorem).
 - (c) If the sectional curvature of a non-compact complete manifold is positive, then it is diffeomorphic to \mathbb{R}^n (Gromoll-Meyer). It is an open conjecture (due to Yau) for Kähler manifolds with $sec > 0$ (for that matter holomorphic bisectional curvature > 0) to be biholomorphic to \mathbb{C}^n .
- (2) Ricci curvature : $Ricc(Y, Z) = tr(X \rightarrow R(X, Y)Z)$, i.e., $Ricc_{ab} = R_{bca}^c$. It turns out that $Ricc(X, Y) = Ricc(Y, X)$. This tensor (like the metric) is symmetric. In fact, one can prove that for 1, 2, 3 dimensions, the Ricci tensor determines the full Riemann tensor. (Not beyond three dimensions though.) The Ricci curvature is very important. Here is a beautiful theorem (Bonnet-Myers) that illustrates its importance.

Theorem 2.2. *If $k > 0$, $m = \dim(M)$, and g is a complete Riemannian metric satisfying $Ric(p) - (m-1)kg(p) \geq 0 \forall p \in M$ (as semi-positive definite matrices), then $\text{diam}(M) \leq \frac{\pi}{\sqrt{k}}$. Hence M is compact. Moreover, since the universal cover is also compact (by pulling back the metric), the fundamental group is finite.*

- (3) Scalar curvature : $S = Ricc_{ab}g^{ab}$. This curvature has the advantage (and disadvantage) of being a single number. The scalar curvature determines the full curvature in dimensions 1, 2. (Actually, all curvatures in dim 1 are zero.) The scalar curvature can be interpreted as follows :

$$(2.3) \quad \frac{\text{Vol}(B_\epsilon(p) \subset M)}{\text{Vol}(B_\epsilon(0) \subset \mathbb{R}^m)} = 1 - \frac{S}{6(m+2)}\epsilon^2 + O(\epsilon^4).$$

Therefore, if $S > 0$, then balls in the manifold have smaller volume because they curve more.

The Yamabe problem asks the following : On a compact manifold, given a metric g_0 , is there is a function f so that $g = e^f g_0$ has constant scalar curvature ? (The answer is now known to be “yes”). If the “constant” of the scalar curvature is negative, then it is not incredibly hard to prove the theorem. There are obstructions to finding metrics of positive scalar curvature on manifolds (you can’t always do it). Even if you can find one, the Yamabe problem in the positive case is very hard. Shockingly enough, its proof involves the positive mass theorem of general relativity.

3. DIVERGENCE, STOKES’ THEOREM, AND LAPLACIANS

Suppose $u : M \rightarrow \mathbb{R}$ is a function on a Riemannian manifold (M, g) whose tangent bundle is equipped with the Levi-Civita connection. Then $\nabla u = \frac{\partial u}{\partial x^j} g^{ij} \frac{\partial}{\partial x^i}$ is called the gradient of u with respect to g . It is just dual to du using the metric g . Suppose c is a regular value of u , then $u^{-1}(c)$ is a submanifold of M of dimension $m-1$. The gradient ∇u is normal to this submanifold. Indeed, if \vec{v} is tangent to the submanifold, i.e., $\vec{v} = \frac{d\gamma}{dt}(0)$ where γ is a curve lying on $u^{-1}(c)$, i.e., $u(\gamma(t)) = c$, then $\frac{du}{dt} = 0$, i.e., $0 = \frac{\partial u}{\partial x^i} \frac{\partial \gamma^i}{\partial t} = (\nabla u)^j g_{ij} \frac{\partial \gamma^i}{\partial t}$. Thus ∇u is perpendicular to \vec{v} .

Suppose X is a smooth vector field. Define the divergence of X

Definition 3.1. $\text{div}(X) = \nabla_i X^i = \frac{\partial X^i}{\partial x^i} + \Gamma_{ik}^i X^k$. So in normal coordinates, it is the usual divergence at p . Note that $\text{div}(fX) = X(f) + f\text{div}(X)$.

Theorem 3.2. $\int_M \text{div}(X) \text{vol}_g = \int_{\partial M} i_X \text{vol}_g$ where $i_X \omega(Y_1, Y_2, \dots) = \omega(X, Y_1, Y_2, \dots)$. If \vec{N} is a unit outward pointing normal vector field on the boundary, then $i_X \text{vol}_g = g(X, \vec{N}) d\text{vol}_{g|_{\partial M}}$.

Proof. Choose oriented normal coordinates x^i for g at $p \in M$. Now

$$\begin{aligned} \text{div}(X)(p) \text{vol}_g(p) &= \sum_i \frac{\partial X^i}{\partial x^i}(p) dx^1 \wedge dx^2 \wedge \dots \wedge dx^m(p) = d\left(\sum_i X^i (-1)^{i-1} dx^1 \wedge \dots \wedge dx^{i-1} \wedge dx^{i+1} \wedge \dots\right)(p) \\ (3.1) \qquad \qquad \qquad &= d(i_X \text{vol})(p) \end{aligned}$$

Since the above equation is an equation of globally defined forms at p , it is independent of coordinates chosen. Thus $\text{div}(X) \text{vol}_g = d(i_X \text{vol})$. By the usual Stokes' theorem, $\int_M \text{div}(X) \text{vol}_g = \int_{\partial M} i_X \text{vol}$. Now if $X = g(X, \vec{N}) \vec{N} + Y$, then Y is tangent to the boundary. Choose oriented normal coordinates such that $x^1 = 0$ corresponds to the boundary (hence $\vec{N}(p) = \frac{\partial}{\partial x^1}$ and Y is a linear combination of ∂_i where $i \geq 2$) Then $i_X \text{vol}(p)|_{x^1=0} = g(X, \vec{N})(p) i_{\vec{N}(p)} dx^1 \wedge dx^2 \wedge \dots(p) + i_{Y(p)} dx^1 \wedge dx^2 \wedge \dots(p) = g(X, \vec{N}) \text{vol}_{g|_{\partial M}}(p)$. As before, this equation holds globally. \square

In particular, if M has no boundary, then $\int_M \text{div}(X) = 0$. Now define

Definition 3.3. The Laplacian Δu where u is a function on M is a function $\Delta u = \text{div}(\nabla u) = \frac{\partial}{\partial x^i} \left(g^{ij} \frac{\partial u}{\partial x^j} \right) + \Gamma_{ik}^i \frac{\partial u}{\partial x^k}$. So in normal coordinates, it is the usual Laplacian at p .

As an example, take the flat metric $g = d\theta^1 \otimes d\theta^1 + d\theta^2 \otimes d\theta^2 + \dots$ on the torus. Then the Laplacian is easily seen to be the Laplacian we studied earlier. Here is an observation using Stokes :

$$(3.2) \qquad \qquad \qquad \int_M \Delta u = \int_M \text{div}(\text{grad}(u)) d\text{vol}_g = 0$$

So if $\Delta u = f$, a necessary condition is that $\int f d\text{vol}_g = 0$ (just like the torus). If $\Delta u = f$, then observe that for any smooth function v ,

$$(3.3) \qquad \int_M v \Delta u \text{vol}_g = \int_M v f \text{vol}_g \Rightarrow \int_M (\text{div}(v \nabla u) - \nabla v \cdot \nabla u) \text{vol}_g = - \int_M \nabla v \cdot \nabla u \text{vol}_g = \int_M u \Delta v \text{vol}_g$$

So we can define a distributional solution of $\Delta u = f$ as an L^2 function u such that the above holds for all smooth v .

What about the curl of a vector field X ? Firstly, given a vector field X , we can produce its dual 1-form $\omega_X(Y) = g(X, Y)$. We can then define $d\omega_X$ as a 2-form. If there is a way to take a 2-form α to an $m-2$ form $*\alpha$, then in 3-dimensions, $*\alpha$ will be a 1-form, whose dual is a vector field. This should be the curl. So we need a notion called the Hodge star $*$ taking k -forms to $m-k$ forms.

Definition 3.4. Given a k -form α on a compact oriented m -dimensional Riemannian manifold (M, g) , $*\alpha$ is a $(m-k)$ -form such that $\alpha \wedge *\beta = \langle \alpha, \beta \rangle_g \text{vol}_g$. Here the inner product on forms is defined as follows : Suppose at p , normal coordinates are chosen, i.e., $g_{ij}(p) = \delta_{ij}$, then $dx^{i_1}(p) \wedge dx^{i_2} \wedge \dots \wedge dx^{i_k}(p)$ form an orthonormal basis at p for k -forms. Note that $\text{vol}(p) = dx^1(p) \wedge dx^2(p) \wedge \dots \wedge dx^m(p)$.